

GRB 030226 in a Density-Jump Medium

Z. G. Dai and X. F. Wu

Department of Astronomy, Nanjing University, Nanjing 210093, China

daizigao@public1.ptt.js.cn, xfwu@nju.edu.cn

ABSTRACT

We present an explanation for the unusual temporal feature of the GRB 030226 afterglow. The R-band afterglow of this burst faded as $\sim t^{-1.2}$ in ~ 0.2 days after the burst, rebrightened during the period of $\sim 0.2 - 0.5$ days, and then declined with $\sim t^{-2.0}$. To fit such a light curve, we consider an ultrarelativistic jetted blast wave expanding in a density-jump medium. The interaction of the blast wave with a large density jump produces relativistic reverse and forward shocks. In this model, the observed rebrightening is due to emissions from these newly forming shocks, and the late-time afterglow is caused by sideways expansion of the jet. Our fitting implies that the progenitor star of GRB 030226 could have produced a stellar wind with a large density jump prior to the GRB onset.

Subject headings: gamma-rays: bursts — relativity — shock waves

1. Introduction

The gamma-ray burst (GRB) 030226 was detected by the *High Energy Transient Explorer 2* Satellite (*HETE-2*) at 03:46:31.99 UT on 26 February 2003 (Suzuki et al. 2003). The burst duration in the 30-400 keV band was beyond 100 seconds, and the fluence in this energy band was $\sim 5.7 \times 10^{-6}$ ergs cm $^{-2}$. Owing to the rapid localization of *HETE-2*, an early afterglow of this burst was observed. Fox, Chen & Price (2003) detected an optical counterpart ~ 0.11 days after the burst, and Price, Fox & Chen (2003) found about 9 minutes later that it had faded. The subsequent spectroscopic observations revealed some absorption lines, indicating that this burst was at a redshift of $z = 1.986 \pm 0.001$ (Greiner et al. 2003a). Thus, the isotropic γ -ray energy release is estimated to be $\sim 6 \times 10^{52}$ ergs, assuming a cosmological model with $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$ and $H_0 = 65$ km s $^{-1}$ Mpc $^{-1}$.

The afterglow light curve of GRB 030226 has one unusual feature. The R-band afterglow declined as $\sim t^{-1.2}$ shown by the four observations during the period of $\sim 0.13 - 0.20$ days after the burst, and about seven hours later, it began to decay with another simple power-law of $t^{-2.0}$ as late-time afterglows from jets usually do. However, contrary to jetted afterglows observed previously, the extrapolation of the early afterglow light curve of GRB 030226 crosses with the

late-time light curve at ~ 2 days rather than ~ 0.5 days, which implies an obvious rebrightening. Such a feature could result from post-burst energy injection. A few variants of this model have been studied (e.g., Dai & Lu 1998a; Panaitescu, Mészáros & Rees 1998; Zhang & Mészáros 2002; Nakar, Piran & Granot 2003). In this Letter, we explain the GRB 030226 afterglow light curve by assuming this burst to be in a density-jump medium (see §2). In §3 we discuss scenarios for density-jump formation and give order-of-magnitude estimates of the density-jump factor, which are consistent with required by our fitting. Our results are summarized in §4.

2. The Model

Before proposing our model, we first discuss implications of the observations on the GRB 030226 afterglow: The observed light curve index (α_2) after ~ 0.5 days since the *HETE-2* trigger suggests that this afterglow might have originated from a highly collimated relativistic shock, because the transition of this shock from the spherical to spreading phase could lead to steepening of the light curve to the flux proportional to t^{-p} for $p > 2$, where p is the spectral index of the shock-accelerated electrons (Rhoads 1999; Sari, Piran & Halpern 1999; for the case of $1 < p < 2$ see Dai & Cheng 2001). Thus, $p = \alpha_2 \simeq 2$ for GRB 030226. If the early afterglow is assumed to result from the shock expanding in a medium with density of $n_w \propto R^{-k}$, then the light curve index in the slow-cooling regime becomes $\alpha_1 = 3(p-1)/4 + k/(8-2k)$ (Dai & Lu 1998b; Mészáros, Rees & Wijers 1998). The observed value of $\alpha_1 \simeq 1.2$, combined with the inferred value of p , implies $k \simeq 2$, revealing a wind surrounding the burst. This environment as an expected signature of massive progenitor stars of GRBs was discussed theoretically in detail (Dai & Lu 1998b; Mészáros et al. 1998; Chevalier & Li 1999, 2000) and confirmed observationally for some bursts (see, e.g., Price et al. 2002 for GRB 011121). Recently, the wind interaction model was also shown to explain well the early afterglow of GRB 021004 (Li & Chevalier 2003).

Therefore, we consider an ultrarelativistic jet with an isotropic energy of E , which first expands in a circumburst wind of the density profile of $n_w(R) = AR^{-2}$ with $A = 3 \times 10^{35} A_* \text{ cm}^{-1}$ for $R < R_0$. Here $A_* = (\dot{M}/10^{-5} M_\odot \text{ yr}^{-1})/(V_w/10^3 \text{ km s}^{-1})$, where $\dot{M}$ and V_w are the mass loss rate and velocity of the wind respectively. After the internal shock emission, the jet will start to sweep up its ambient wind, leading to an ultrarelativistic blast wave. The initial hydrodynamics and emission were recently discussed by Wu et al. (2003) and Kobayashi & Zhang (2003). We further assume this blast wave which will hit an outer high-density homogeneous region for $R \geq R_0$. Dai & Lu (2002, hereafter DL02) have analyzed the hydrodynamics and afterglow emission from a relativistic blast wave in detail when it expands in a density-jump medium, which might have been produced around $\sim 10^{18} \text{ cm}$ due to the interaction of a stellar wind with the ambient matter (see §3).

The interaction of a relativistic blast wave with a large density jump can be described through two shocks: a reverse shock that propagates into the hot shell (viz., the $R \leq R_0$ wind matter swept up by the blast wave), and a forward shock that propagates into the outer high-density

medium. Thus, there are four regions separated in the system of interest by the two shocks: (1) the unshocked high-density medium, (2) the forward-shocked high-density medium, (3) the reverse-shocked hot shell, and (4) the unshocked hot shell. We denote γ_i as the Lorentz factor of region “ i ” measured in the local medium’s rest frame, and n_1 is the baryon number density of the medium at $R \geq R_0$ (region 1). Defining the ratio (f) of the energy densities of regions 4 and 1, DL02 have got

$$\gamma_2 = \gamma_3 = \frac{\gamma_4^{1/2} f^{1/4}}{3^{1/4}}, \quad (1)$$

and found that a density jump with a factor of much more than 21 can lead to a relativistic reverse shock. In this case, the Lorentz factor of regions 2 and 3 can further be scaled as $\gamma_2 = \gamma_3 = 1.4(1+z)^{1/2} A_*^{1/4} n_{1,3}^{-1/4} t_0^{-1/2} (R/R_0)^{-1}$, where $n_{1,3} = n_1/10^3 \text{ cm}^{-3}$, and t_0 is the observed time (in 1 day) at $R = R_0 = 8.0 \times 10^{17} (1+z)^{-1/2} E_{53}^{1/2} A_*^{-1/2} t_0^{1/2} \text{ cm}$ (where E_{53} is the isotropic-equivalent energy of the jet in units of 10^{53} ergs). Comparing γ_2 (or γ_3) with the Lorentz factor of region 4 at the crossing time, $\gamma_4 = 8.8(1+z)^{1/4} E_{53}^{1/4} A_*^{-1/4} t_0^{-1/4}$, it is easy to see that the Lorentz factor of region 3 is much less than that of region 4 for typical parameters, showing that most of the initial kinetic energy is converted into thermal energy by the shocks. The radius at which the reverse shock has just crossed region 4 is derived as $R_\Delta = R_0[1 + 1.25 \times 10^{-2} (1+z)^{1/2} E_{53}^{-1/2} A_* n_{1,3}^{-1/2} t_0^{-1/2}]$ (DL02). Thus, $R_\Delta/R_0 - 1 \ll 1$ for typical parameters, so that the Lorentz factor of region 2 at $R = R_\Delta$ is well approximated by its Lorentz factor at $R = R_0$. If the initial half opening angle of the jet is $\theta_0 \sim 0.1$ as in most of the afterglows analyzed by Frail et al. (2001), then it is possible that after the crossing time the Lorentz factor of the jet is far below θ_0^{-1} and therefore the flux decays as t^{-p} due to sideways expansion as long as the jet is still relativistic (Rhoads 1999; Sari et al. 1999).

In the standard afterglow model (Sari, Piran & Narayan 1998), the synchrotron flux $F_{\nu,i}$ of region “ i ” in optical to X-ray bands is determined by three quantities: the typical synchrotron frequency $\nu_{m,i}$, the cooling frequency $\nu_{c,i}$, and the peak flux $F_{\nu,max,i}$. We divide the whole evolution into three stages: (I) before, (II) during, and (III) after the crossing time. Using Chevalier & Li’s (2000) results, we get these three quantities at stage I: $\nu_m^I(t) = 4.8 \times 10^{13} (1+z)^{1/2} \epsilon_{e,-1}^2 \epsilon_{B,-1}^{1/2} \xi^2 E_{53}^{1/2} t^{-3/2} \text{ Hz}$, $\nu_c^I(t) = 6.6 \times 10^{13} (1+z)^{-3/2} \epsilon_{B,-1}^{-3/2} E_{53}^{1/2} A_*^{-2} t^{1/2} \text{ Hz}$, and $F_{\nu,max}^I(t) = 0.59 (1+z)^{3/2} \epsilon_{B,-1}^{1/2} E_{53}^{1/2} A_* t^{-1/2} D_{L,28}^{-2} \text{ Jy}$, where $\xi = (p-2)/(p-1)$, $\epsilon_e = 0.1 \epsilon_{e,-1}$ and $\epsilon_B = 0.1 \epsilon_{B,-1}$ are constant fractions of the internal energy density going into the electrons and the magnetic fields respectively, t is the observer’s time in units of 1 day, and $D_{L,28}$ is the luminosity distance to the source in units of 10^{28} cm .

At stage II, the spectral break frequencies and the peak flux evolve as

$$\nu_{m,i}^{\text{II}} = \begin{cases} 1.3 \times 10^{12} (1+z) \epsilon_{e,-1}^2 \epsilon_{B,-1}^{1/2} \xi^2 A_* n_{1,3}^{-1/2} t_0^{-2} (R/R_0)^{-4} \text{ Hz}, & \text{if } i = 2, \\ 1.0 \times 10^{15} \epsilon_{e,-1}^2 \epsilon_{B,-1}^{1/2} \xi^2 E_{53} A_*^{-1} n_{1,3}^{1/2} t_0^{-1} (R/R_0)^{-2} \text{ Hz}, & \text{if } i = 3, \\ \nu_m^I(t_0) (R/R_0)^{-4}, & \text{if } i = 4. \end{cases} \quad (2)$$

$$\nu_{c,i}^{\text{II}} = \begin{cases} 1.0 \times 10^{12} (1+z)^{-1} \epsilon_{B,-1}^{-3/2} A_*^{-1} n_{1,3}^{-1/2} (t/t_0)^{-2} (R/R_0)^4 \text{Hz}, & \text{if } i = 2, 3, \\ \nu_c^{\text{I}}(t_0) (R/R_0)^{-4}, & \text{if } i = 4. \end{cases} \quad (3)$$

$$F_{\nu,max,i}^{\text{II}} = \begin{cases} 470 (1+z)^{1/2} \epsilon_{B,-1}^{1/2} E_{53}^{1/2} A_*^{-1} n_{1,3}^{1/2} D_{L,28}^{-2} [(R/R_0)^3 - 1] (R/R_0)^{-2} \text{Jy}, & \text{if } i = 2, \\ 28 (1+z)^{1/2} \epsilon_{B,-1}^{1/2} E_{53}^{1/2} n_{1,3}^{1/2} D_{L,28}^{-2} [(R/R_0)^2 - 1] (R/R_0)^{-2} \text{Jy}, & \text{if } i = 3, \\ F_{\nu,max}^{\text{I}}(t_0) \zeta (R/R_0)^{-2}, & \text{if } i = 4, \end{cases} \quad (4)$$

where ζ is the ratio of the total electron numbers of region 4 at radius $R \in (R_0, R_\Delta)$ and at radius R_0 . It should be emphasized that $\nu_{c,4}^{\text{II}}$ in equation (3) is in fact a cutoff frequency because of adiabatic expansion of region 4, implying that there is no emission from this region at frequencies above $\nu_{c,4}^{\text{II}}$.

From DL02, we obtain the time at which $R = R_\Delta$ as follows: $t_{\Delta,2} = t_{\Delta,3} = 2.0t_0$ for regions 2 and 3. If the afterglow originates from a jet, one expects that at stage III the break frequencies and the peak flux of region 2 are given by $\nu_{m,2}^{\text{III}} = \nu_{m,2}^{\text{II}}(t/t_{\Delta,2})^{-2}$, $\nu_{c,2}^{\text{III}} = \nu_{c,2}^{\text{II}}(t/t_{\Delta,2})$, and $F_{\nu,max,2}^{\text{III}} = F_{\nu,max,2}^{\text{II}}(t/t_{\Delta,2})^{-1}$ (Sari et al. 1999). For the parameters adopted below, the cutoff frequency of region 3 is less than the optical frequency so that there is no emission from this region at frequencies above the optical band.

Figure 1 presents an example case of our fitting to the multiwavelength light curves of the GRB 030226 afterglow. Please note that an error of 0.05 mag for the initial four afterglow data points in the figure has been taken as the error in the observed brightness of two calibration stars by Garnavich, von Braun & Stanek (2003). This error is smaller than a conservative estimate of 0.3 mag by Zeh et al. (2003). The model parameters are taken: $\epsilon_{e,-1} = 0.9$, $\epsilon_{B,-1} = 4$, $E_{53} = 1.0$, $A_* = 0.01$, $p = 2.01$, $t_0 = 0.22$ days, and $n_{1,3} = 0.1$. If the emission off the line of sight (LOS) were neglected, one would see an initial drop at $t \simeq t_0$, a rapid brightening at $t_0 \lesssim t \lesssim t_{\Delta,2}$ and an abrupt decay at $t \simeq t_{\Delta,2}$ for all the light curves (DL02). However, such features are expected to be smoothed by the off-LOS emission. In Figure 1, we have considered the off-LOS emission effect for region 4 due to its ultrarelativistic expansion (e.g., $\gamma_4 \sim 53 \gg 1$ for the taken model parameters) at $t \simeq t_0$, following Kumar & Panaitescu (2000). However, this effect is insignificant for region 3 because $\gamma_3 \sim 2.9$ is not far greater than unity at $t \simeq t_{\Delta,2}$.

We next discuss implications of the fitting. First, because the late-time afterglow at high frequencies in our model results from fast-cooling electrons, the spectral index should be $\beta = p/2 \simeq 1.0$, which is consistent with the observed values, $\beta_{\text{opt,ob}} = 0.99 \pm 0.42$ at the optical band (Nysewander et al. 2003) and $\beta_{\text{X,ob}} = 1.0 \pm 0.1$ at the X-ray band (Pedersen et al. 2003; Sako & Fox 2003), respectively. Second, the jet's isotropic-equivalent energy ($\sim 10^{53}$ ergs) from our fitting is close to the isotropic γ -ray energy release ($\sim 6 \times 10^{52}$ ergs). This shows that the efficiency of γ -ray emission from internal shocks is as high as $\sim 40\%$. Third, our fitting gives a weak wind $A_* \sim 0.01$, which implies non-detection of an optical flash from GRB 030226, as in most of the GRBs observed previously (Wu et al. 2003). Finally, the required value of n_1 indicates that the progenitor star of GRB 030226 could have been in a dense medium such as a giant molecular cloud (GMC).

3. Density-Jump Formation

In this section we show that a large density jump seems to form in the vicinity of a massive progenitor star. A natural scenario for density-jump formation is the interaction of a wind from a massive star with its outer environment. This interaction produces a stellar wind bubble. Numerical simulations of the bubble dynamics in a homogeneous interstellar medium were performed by Ramirez-Ruiz et al. (2001), and here we give analytical studies to obtain the order-of-magnitude factor of a density jump by considering two kinds of environment. First, we discuss a simple case in which the outer environment of a massive star is a homogeneous GMC with density n_1 . Owing to the presence of two shocks (viz., a reverse shock that propagates into the wind gas and a forward shock that propagates into the GMC gas), the system has a four-zone structure consisting of (1) the unshocked GMC gas, (2) the shocked GMC shell, (3) the shocked wind gas, and (4) the unshocked wind gas moving with a velocity V_w and with a mass loss rate $\dot{M}$, where zones 2 and 3 are separated by a contact discontinuity. Assuming an adiabatic expansion and two strong shocks (Castor, McCray & Weaver 1975; Wijers 2001; Mirabal et al. 2003), we get the forward shock radius

$$R_{\text{sh}}^f \simeq 4.8 \times 10^{18} \dot{M}_{-7}^{1/5} V_{w,3}^{2/5} n_{1,2}^{-1/5} t_{w,5}^{3/5} \text{ cm}, \quad (5)$$

and the reverse shock radius

$$R_{\text{sh}}^r \simeq 7.0 \times 10^{17} \dot{M}_{-7}^{3/10} V_{w,3}^{1/10} n_{1,2}^{-3/10} t_{w,5}^{2/5} \text{ cm}, \quad (6)$$

where $\dot{M}_{-7} = \dot{M}/10^{-7} M_{\odot} \text{ yr}^{-1}$, $V_{w,3} = V_w/10^3 \text{ km s}^{-1}$, $n_{1,2} = n_1/10^2 \text{ cm}^{-3}$, and $t_{w,5}$ is the wind lifetime in units of 10^5 years. Here the reason for scaling the stellar mass loss rate with $10^{-7} M_{\odot} \text{ yr}^{-1}$ is that $A_* \sim 0.01$ in our fitting implies $\dot{M} \sim 10^{-7} V_{w,3} M_{\odot} \text{ yr}^{-1}$. This scaling is typical for main-sequence stellar winds, and further not far below typical mass loss rates of Wolf-Rayet stars (Willis 1991). One large density jump appears at the contact discontinuity, whose factor is approximated by

$$\chi = \frac{n_c}{n_w(R_{\text{sh}}^r)} \sim 1.7 \times 10^4 \dot{M}_{-7}^{-2/5} V_{w,3}^{6/5} n_{1,2}^{2/5} t_{w,5}^{4/5}. \quad (7)$$

The radius at which the density jump is required to appear in our fitting is estimated as $R_0 \sim 2.2 \times 10^{18} \text{ cm}$, which is larger than R_{sh}^r and closer to R_{sh}^f for reasonable wind parameters, so the afterglow shock wave would be able to reach the density jump at the early time when the rebrightening was observed in the GRB 030226 afterglow light curve.

Second, the surrounding medium of a fast wind from a massive star (e.g., a Wolf-Rayet star) may be a dense slow wind, because this star could have passed through a red supergiant (RSG) phase. During such a phase, the star has produced a wind with a larger mass loss rate $\dot{M}_{\text{RSG}}$ and a slower velocity V_{RSG} . This wind has formed a medium with density $n_{\text{RSG}}(R) = 3 \times 10^{37} (\dot{M}_{\text{RSG},-5}/V_{\text{RSG},1}) R^{-2} \text{ cm}^{-3}$, where $\dot{M}_{\text{RSG},-5} = \dot{M}_{\text{RSG}}/10^{-5} M_{\odot} \text{ yr}^{-1}$ and $V_{\text{RSG},1} = V_{\text{RSG}}/10 \text{ km s}^{-1}$. The interaction of a RSG wind with its homogeneous environment (e.g., a GMC) could have given rise to a RSG wind bubble. The dynamics of this bubble is similar to the case discussed above, provided that they are both adiabatic. From

equation (5), we can write directly the radius of a thin shell swept up by the RSG wind as $R_{\text{sh,RSG}} \simeq 1.9 \times 10^{18} \dot{M}_{\text{RSG},-5}^{1/5} V_{\text{RSG},1}^{2/5} n_{1,2}^{-1/5} t_{\text{RSG},5}^{3/5}$ cm, where $t_{\text{RSG},5}$ is the RSG wind lifetime in units of 10^5 years. Also, we obtain the factor of a density jump appearing at the contact discontinuity in the RSG wind bubble, $\chi_{\text{RSG}} \sim 11 \dot{M}_{\text{RSG},-5}^{-2/5} V_{\text{RSG},1}^{6/5} n_{1,2}^{2/5} t_{\text{RSG},5}^{4/5}$.

Furthermore, a fast wind from the Wolf-Rayet phase sweeps up the pre-existing slow wind, leading possibly to another wind bubble. Chevalier & Imamura (1983) gave self-similar solutions of this bubble. Assuming an adiabatic evolution and two strong shocks (Garcia-Segura & Mac Low 1995), we have the forward shock radius

$$R_{\text{sh}}^{\text{f}} \simeq 8.6 \times 10^{18} \dot{M}_{-7}^{1/3} V_{w,3}^{2/3} \dot{M}_{\text{RSG},-5}^{-1/3} V_{\text{RSG},1}^{1/3} t_{w,5} \text{ cm}, \quad (8)$$

and the reverse shock radius

$$R_{\text{sh}}^{\text{r}} \simeq 1.6 \times 10^{18} \dot{M}_{-7}^{1/2} V_{w,3}^{1/2} \dot{M}_{\text{RSG},-5}^{-1/2} V_{\text{RSG},1}^{1/2} t_{w,5} \text{ cm}. \quad (9)$$

Letting R_{sh}^{f} in equation (8) be equal to $R_{\text{sh,RSG}}$, we define a critical lifetime of the Wolf-Rayet wind,

$$t_w^{\text{cr}} \simeq 2.2 \times 10^4 \dot{M}_{-7}^{-1/3} V_{w,3}^{-2/3} \dot{M}_{\text{RSG},-5}^{8/15} V_{\text{RSG},1}^{1/15} n_{1,2}^{-1/5} t_{\text{RSG},5}^{3/5} \text{ yrs}. \quad (10)$$

If $t_w \lesssim t_w^{\text{cr}}$, the Wolf-Rayet wind only sweeps up the RSG wind. In this subcase, a large density jump occurs at the contact discontinuity and its factor is $\chi_w = n_{\text{RSG}}(R_{\text{sh}}^{\text{f}})/n_w(R_{\text{sh}}^{\text{r}}) \sim 360 \dot{M}_{-7}^{-2/3} V_{w,3}^{2/3} \dot{M}_{\text{RSG},-5}^{2/3} V_{\text{RSG},1}^{-2/3}$, where R_{sh}^{f} and R_{sh}^{r} are given by equations (8) and (9) respectively. When $R_{\text{sh}}^{\text{f}} \sim R_{\text{sh,RSG}}$, the above two density jumps merge to one larger density jump with a factor of

$$\chi \sim \chi_w \chi_{\text{RSG}} \sim 4 \times 10^3 \dot{M}_{-7}^{-2/3} V_{w,3}^{2/3} \dot{M}_{\text{RSG},-5}^{4/15} V_{\text{RSG},1}^{8/15} n_{1,2}^{2/5} t_{\text{RSG},5}^{4/5}. \quad (11)$$

On the other hand, if $t_w > t_w^{\text{cr}}$, the Wolf-Rayet wind not only sweeps up the RSG wind but also the outer GMC gas, in which subcase the dynamics can be approximated by equations (5) and (6), and thus the factor of a large density jump that occurs at the contact discontinuity is estimated by equation (7). Because R_0 is larger than R_{sh}^{r} and closer to R_{sh}^{f} in both subcases, the afterglow shock wave would be able to arrive at the large density jump at an expected early time of GRB 030226.

4. Conclusions

In this paper we have shown that density jumps (or bumps) are likely to occur in the vicinity of a GRB, especially if a stellar wind has interacted with the ambient matter or if winds ejected at different velocities have collided each other, prior to the GRB onset. We have considered an ultrarelativistic *jetted* blast wave in a density-jump medium to explain the unusual temporal feature of the GRB 030226 afterglow. A large density jump can produce a relativistic reverse shock, which leads naturally to (1) effective conversion of the initial kinetic energy to thermal energy and (2) rapid deceleration of the jet. Therefore, the assumption of one large density jump

has two advantages. Based on them, the observed rebrightening is understood as due to emissions from reverse and forward shocks forming in the interaction of the blast wave with the density jump, and the late-time afterglow is caused by sideways expansion of the jet.

We thank the referee and X. Y. Wang for valuable comments. This work was supported by the National Natural Science Foundation of China (grant 10233010) and the National 973 Project (NKBRF G19990754).

REFERENCES

- Ando, M. et al. 2003a, GCN Circ. 1882
- Ando, M. et al. 2003b, GCN Circ. 1884
- Castor, J., McCray, R., & Weaver, R. 1975, ApJ, 200, L107
- Chevalier, R. A., & Imamura, J. N. 1983, ApJ, 270, 554
- Chevalier, R. A., & Li, Z. Y. 1999, ApJ, 520, L29
- Chevalier, R. A., & Li, Z. Y. 2000, ApJ, 536, 195
- Covino, S. et al. 2003, GCN Circ. 1909
- Dai, Z. G., & Cheng, K. S. 2001, ApJ, 558, L109
- Dai, Z. G., & Lu, T. 1998a, Phys. Rev. Lett., 81, 4301
- Dai, Z. G., & Lu, T. 1998b, MNRAS, 298, 87
- Dai, Z. G., & Lu, T. 2002, ApJ, 565, L87 (DL02)
- Fatkhullin, T. et al. 2003, GCN Circ. 1925
- Fox, D. W., Chen, H. W., & Price, P. A. 2003, GCN Circ. 1879
- Frail, D. A. et al. 2001, ApJ, 562, L55
- Garcia-Segura, G. G., & Mac Low, M. M. 1995, ApJ, 455, 145
- Garnavich, P., von Braun, K., & Stanek, K. 2003, GCN Circ. 1885
- Greiner, J., Guenther, E., Klose, S., & Schwarz, R. 2003a, GCN Circ. 1886
- Greiner, J., Ries, C., Barwig, H., Fynbo, J., & Klose, S. 2003b, GCN Circ. 1894
- Guarnieri, A. et al. 2003, GCN Circ. 1892

- Kobayashi, S., & Zhang, B. 2003, ApJL, submitted (astro-ph/0304086)
- Kumar, P., & Panaitescu, A. 2000, ApJ, 541, L51
- Li, Z. Y., & Chevalier, R. A. 2003, ApJ, 589, L69
- Maiorano, E. et al. 2003, GCN Circ. 1933
- Mészáros, P., Rees, M. J., & Wijers, R. A. M. J. 1998, ApJ, 499, 301
- Mirabal, N. et al. 2003, ApJ, submitted (astro-ph/0303616)
- Nakar, E., Piran, T., & Granot, J. 2003, NewA, 8, 495
- Nysewander, M. C., Moran, J., Reichart, D., Henden, A., & Schwartz, M. 2003, GCN Circ. 1921
- Panaitescu, P., Mészáros, P., & Rees, M. J. 1998, ApJ, 503, 314
- Pedersen, K., Fynbo, J., Hjorth, J., & Watson, D. 2003, GCN Circ. 1924
- Price, P. A. et al. 2002, ApJ, 572, L51
- Price, P. A., Fox, D. W., & Chen, H. W. 2003, GCN Circ. 1880
- Price, P. A., & Warren, B. E. 2003, GCN Circ. 1890
- Ramirez-Ruiz, E., Dray, L. M., Madau, P., & Tout, C. A. 2001, MNRAS, 327, 829
- Rhoads, J. 1999, ApJ, 525, 737
- Rumyantsev, V., Biryukov, V., & Pozanenko, A. 2003, GCN Circ. 1908
- Rumyantsev, V., Sergeeva, L., & Pozanenko, A. 2003, GCN Circ. 1929
- Sako, M., & Fox, D. W. 2003, GCN Circ. 1928
- Sari, R., Piran, T., & Halpern, J. P. 1999, ApJ, 519, L17
- Sari, R., Piran, T., & Narayan, R. 1998, ApJ, 497, L17
- Semkov, E. 2003, GCN Circ. 1935
- Suzuki, M. et al. 2003, GCN Circ. 1888
- von Braun, K., Garnavich, P., & Stanek, K. 2003a, GCN Circ. 1881
- von Braun, K., Garnavich, P., & Stanek, K. 2003b, GCN Circ. 1902
- Wijers, R. A. M. J. 2001, in Gamma-Ray Bursts in the Afterglow Era, eds. E. Costa, F. Frontera & J. Hjorth (Springer), 306

- Willis, A. J. 1991, in *Wolf-Rayet Stars and Interrelations with Other Massive Stars in Galaxies*, eds. K. A. van der Hulcht & B. Hidayat (Dordrecht: Reidel), 265
- Wu, X. F., Dai, Z. G., Huang, Y. F., & Lu, T. 2003, *MNRAS*, in press (astro-ph/0304110)
- Zeh, A., Klose, S., Greiner, J., Fynbo, J., & Jakobsson, P. 2003, *GCN Circ.* 1898
- Zhang, B., & Mészáros, P. 2002, *ApJ*, 566, 712




Fig. 1.— Fitting of the multiwavelength afterglow of GRB 030226 by assuming an ultrarelativistic blast wave which expands in a density-jump medium. The model parameters are seen in the text. Data are from Ando et al. 2003a, 2003b; Covino et al. 2003; Fatkhullin et al. 2003; Garnavich, von Braun & Stanek 2003; Greiner et al. 2003b;Guarnieri et al. 2003; Maiorano et al. 2003; Nysewander et al. 2003; Price & Warren 2003; Rummyantsev, Biryukov, & Pozanenko 2003; Rummyantsev, Sergeeva & Pozanenko 2003; Semkov 2003; von Braun, Garnavich & Stanek 2003a, 2003b; Zeh et al. 2003.